



COVID-19 SCREENING FLOW CHARTS

Chippewas of Kettle & Stony Point First Nation



JANUARY 2022

You have symptoms and are concerned you may have COVID-19. Now What?

Do you have any of these symptoms: Fever / chills, cough, shortness of breath, decrease / loss of smell and taste?

No

Yes

- Sore throat
- Headache

Do you have two or more of these symptoms?:

- Extreme fatigue
- Runny nose / nasal congestion
- Muscle aches / joint pain
- GI Symptoms (i.e, vomiting or diarrhea)

No

Yes

- It is less likely that you have COVID-19 infection.
- Self-isolate until your symptoms are improving for at least 24 hours (48 hours for gastrointestinal symptoms.)
- Your household members do not need to self-isolate.

- It is highly likely that you have a COVID-19 infection.
- You must **self-isolate** immediately and seek COVID-19 Testing (Kettle and Stony Point Health Services).
- **Isolation period will be at least 10 days from symptom onset** and until symptoms are improving for 24 hours (or 48 hours for gastrointestinal symptoms).
- All of your household members (regardless of their vaccination status) must self-isolate while you are self-isolating. If household members become symptomatic, they should also seek testing.
- If your symptoms worsen, seek advice from Telehealth, KSPHS or your health care provider.
- Notify your workplace.

Note: In the context of Omicron, individuals who are previously positive in the last 90 days and not fully vaccinated are not considered equivalent to fully vaccinated.




COVID-19 SCREENING FLOW CHARTS

Chippewas of Kettle & Stony Point First Nation



JANUARY 2022

You've been exposed to someone who has tested positive for COVID-19 on PCR, rapid molecular, or rapid antigen test. Now What?



Note: In the context of Omicron, individuals who are previously positive in the last 90 days and not fully vaccinated are not considered equivalent to fully vaccinated.